



TC-89 PRIMER

PRODUCT DESCRIPTION:

This primer was developed to apply against cured urethane substrates for achieving good adhesion when applying another urethane product for painting or bonding applications.

APPLICATION:

Apply to clean and dry surface to 1/10 mil or less in thickness. Spraying is the preferred method of applying this primer, however a light brush or roller coat is adequate in some cases. Drying times will vary with working conditions. At room temperature, the primer is generally tack free in 45 minutes to 1 hour.

It is recommended that best adhesion results will be gained by applying coatings, casting materials or other materials on the primer between 45 and 60 minutes after the primer has been applied. Mixed TC-89 A/B should be used within 1 week.

COVERAGE:

1 pint will cover approximately 40 square feet and will provide about 1/2 mil dry thickness. NOTE: This may vary depending on the substrate's absorption characteristics.

STORAGE:

Store in a cool dry place. Unopened containers will have a shelf life of 6 months from date of shipment when properly stored at room temperatures. Purge opened containers with dry nitrogen before re-sealing.

HANDLING PROPERTIES:

Mix Ratio (by weight):

- Part A 10 parts by weight
- Part B 100 parts by weight

Specific Gravity:

- Part A 1.23
- Part B 0.81

Color:

- Part A Dark Brown
- Part B Clear

SAFETY PRECAUTIONS:

Use in a well-ventilated area. Avoid contact with skin using protective gloves and protective clothing. Repeated or prolonged contact on the skin may cause an allergic reaction.

Eye protection is extremely important. Always use approved safety glasses or goggles when handling this product.

IF CONTACT OCCURS:

Skin: Immediately wash with soap and water. Remove contaminated clothing and launder before reuse. It is *not* recommended to remove resin from skin with solvents. Solvents only increase contact and dry skin. Seek qualified medical attention if allergic reactions occur.

Eyes: Immediately flush with water for at least 15 minutes. Call a physician.

Ingestion: If swallowed, call a physician immediately. Remove stomach contents by gastric suction or induce vomiting only as directed by medical personnel. Never give anything by mouth to an unconscious person.

Refer to the Material Safety Data Sheet before using this product.